

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
8 November 2001 (08.11.2001)

PCT

(10) International Publication Number
WO 01/82916 A2(51) International Patent Classification⁷: A61K 31/00

(21) International Application Number: PCT/US01/14393

(22) International Filing Date: 2 May 2001 (02.05.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/201,613 3 May 2000 (03.05.2000) US

(71) Applicant (for all designated States except US): TULARIK INC. [US/US]; Two Corporate Drive, South San Francisco, CA 94080 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): JAEN, Juan, C. [US/US]; 154 Los Robles Drive, Burlingame, CA 94010 (US). CHEN, Jin-Long [CN/US]; 101 Challenge Court, Foster City, CA 94404 (US).

(74) Agents: SNYDER, Joseph, R.; Townsend and Townsend and Crew LLP, Two Embarcadero Center, 8th floor, San Francisco, CA 94111-3834 et al. (US).


(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: COMBINATION THERAPEUTIC COMPOSITIONS AND METHODS OF USE



(57) Abstract: The present invention provides pharmaceutical compositions and methods for the treatment of diabetes mellitus using combination therapy. The compositions relate to a compound of Formula I and an antidiabetic agent such as sulfonylureas, biguanides, glitazones, α -glucosidase inhibitors, potassium channel antagonists, aldose reductase inhibitors, glucagon antagonists, activators of RXR, insulin therapy or other anti-obesity agent. The methods include the administration of the combination of compound of Formula I with antidiabetic agent where the two components are delivered in a simultaneous manner, where the compound of Formula I is administered first, followed by the antidiabetic agent, as well as wherein the antidiabetic agent is delivered first followed by the compound of Formula I.

COMBINATION THERAPEUTIC COMPOSITIONS AND METHODS OF USE

CROSS REFERENCE TO RELATED APPLICATIONS

5 This application claims priority to U.S. Provisional Application No. 60/201,613, filed May 3, 2000, the teaching of which are hereby incorporated by reference for all purposes.

FIELD OF INVENTION

10 In general, the present invention relates to pharmaceutical compositions, and more particularly, to pharmaceutical compositions for the treatment of diabetes mellitus using combination therapy.

BACKGROUND OF THE INVENTION

15 Diabetes mellitus is a term generally used to refer to various pathological states characterized by hyperglycemia and altered metabolism of lipids, carbohydrates and proteins. These conditions are also often associated with other co-morbidities, such as obesity and an increased risk of cardiovascular disease. By some estimates, as many as 600,000 new individuals become clinically diabetic every year in the United States.

20 Diabetic conditions are generally classified as either insulin-dependent diabetes mellitus (IDDM, Type I diabetes) or non-insulin-dependent diabetes mellitus (NIDDM, Type II diabetes). There are also less common clinical pathologies that are associated with diabetic conditions, such as gestational maturity-onset diabetes of youth (MODY), tropical diabetes secondary to chronic pancreatitis, diabetes secondary to
25 pancreatic disease or surgery, and diabetes secondary to endocrinopathies.

 Virtually all forms of diabetes are due to a decrease in the circulating concentration of insulin (insulin deficiency) and/or a decrease in the response of peripheral tissues to insulin (insulin resistance). These abnormalities lead to alterations in the metabolism of carbohydrates, lipids, ketones and amino acids, and a hyperglycemic
30 condition. IDDM appears to have an autoimmune etiology, which results in destruction of β islet cells in the pancreas and the resulting inability to produce insulin. The etiology of NIDDM, the most prevalent form of diabetes, is more complex and possibly heterogeneous. Some loss of β -cell volume is generally noted in these patients; as well as

decreased circulating levels of insulin. NIDDM patients may also suffer commonly from insulin resistance.

The best-established therapy for all IDDM and many NIDDM patients is subcutaneous insulin treatment. Additionally, insulin is used as the treatment of choice for patients with postpancreatectomy diabetes or gestational diabetes. While insulin is a key element in the control of these hyperglycemic conditions, there are a number of limitations associated with its use, including hypoglycemia, allergic reactions to insulin, lipoatrophy, lipohypertrophy, body weight gain, edema, and insulin resistance. There are a number of new forms of insulin on the market or in various stages of clinical evaluation, including new delivery systems, various recombinant forms, new routes of administration, and gene therapy. These novel forms of insulin treatments are believed to share some of the same limitations outlined above. A significant improvement in the treatment of diabetes can be achieved if insulin treatment is combined with agents that increase the insulin sensitivity of the peripheral tissues.

The concept of combination therapy is well exploited in current medical practice. Treatment of a pathology by combining two or more agents that target the same pathogen or biochemical pathway sometimes results in greater efficacy and diminished side effects relative to the use of the therapeutically relevant dose of each agent alone. In some cases, the efficacy of the drug combination is *additive* (the efficacy of the combination is approximately equal to the sum of the effects of each drug alone), but in other cases the effect can be *synergistic* (the efficacy of the combination is greater than the sum of the effects of each drug given alone). In real medical practice, it is often quite difficult to determine if drug combinations are additive or synergistic.

For most diabetic patients, treatment involves some form of insulin therapy. In addition, IDDM patients may receive a biguanide (*e.g.*, metformin) to enhance the insulin utilization by peripheral tissues. NIDDM patients are often treated with a combination of insulin, a sulfonylurea (to enhance insulin production in the pancreas) and a biguanide or glitazone (to enhance insulin sensitivity by peripheral tissues). For example, the improved utility of a glitazone in combination with a sulfonylurea was recently demonstrated in human clinical trials (*see*, WO 98/36755). Recently, two glitazone compounds (rosiglitazone and pioglitazone) were approved in the United States for the treatment of NIDDM patients in combination with metformin.

A variety of antidiabetic compounds are known. For example, sulfonylureas are a group of drugs that induce hypoglycemia by stimulating insulin

release from the pancreas. Generally, sulfonylureas have found wide utility in the treatment of NIDDM. Their efficacy is decreased in IDDM because of the inherent inability of the patient to produce insulin. Adverse reactions to sulfonylureas occur in a fraction of patients, particularly the elderly. One of the most severe side effects is hypoglycemia and coma. Other side effects include nausea and vomiting, cholestatic jaundice, agranulocytosis, cardiovascular mortality, aplastic and hemolytic anemias, generalized hypersensitivity reactions and dermatological reactions.

Biguanides are another group of drugs, first introduced in the mid 1950's, that have shown efficacy in the treatment of hyperglycemia by mechanisms that are not well understood. The best known agents of this type include metformin, phenformin and buformin. Unlike the sulfonylureas, metformin does not induce release of insulin from the pancreas. It is thought that its effects are mediated by increasing insulin activity in peripheral tissues, reducing hepatic glucose output due to inhibition of gluconeogenesis and reducing the absorption of glucose from the intestine. Side effects associated with the use of biguanides include lactic acidosis, diarrhea, nausea, and anorexia. These agents are often given in combination with drugs that increase the output of insulin from the pancreas, such as the sulfonylureas, which sometimes results in greater efficacy and/or the ability to use lower doses of the drugs, with an improved side effect profile.

More recently, the glitazones have been introduced and are widely used in the treatment of NIDDM. These agents, also known generically as thiazolidinediones, such as troglitazone, rosiglitazone and pioglitazone, are thought to work by increasing the sensitivity of peripheral tissues, such as skeletal muscle, towards insulin. They are often used in combination with insulin or other agents, such as the sulfonylureas, that enhance the release of insulin from the pancreas. A number of side effects have been described during the clinical evaluation of these agents, including hepatotoxicity, organomegaly, edema, anemia and body weight gain. While hepatotoxicity may be the most acutely life-threatening of these conditions, it does not appear in a large percentage of the patient population. On the other hand, the increases in body weight gain associated with chronic glitazone treatment are generally perceived as worsening an already critical co-morbid condition in the majority of the diabetic patients, and may ultimately result in the loss of antidiabetic efficacy for this type of agent after chronic treatment.

α -Glucosidase inhibitors, such as acarbose, reduce intestinal absorption of starch, dextrin, and disaccharides by inhibiting the action of intestinal brush border α -

glucosidase. Inhibition of this enzyme slows the absorption of carbohydrates and the rise in plasma glucose that normally follows after a meal is blunted. Acarbose has shown some benefit in IDDM and NIDDM patients, but is often associated with dose-related malabsorption, flatulence and abdominal bloating.


Other types of agents that have found limited utility in treating diabetes include potassium channel antagonists such as repaglinide, and aldose reductase inhibitors such as zopolrestat and tolrestat. Still in the experimental stage, glucagon antagonists, activators of the retinoid-X receptor (RXR), activators of PPAR α , activators of PPAR δ and anti-obesity agents are also being evaluated as potential antidiabetic agents.

In view of the foregoing, there remains a need in the art to provide more efficacious treatment for diabetic conditions and diabetic complications. Combination therapy treatments are needed that will reduce the amount of drugs taken, thereby decreasing side effects. The present invention fulfills these and other needs.

SUMMARY OF THE INVENTION

The present invention provides pharmaceutical compositions for the treatment of a variety of diseases, including diabetes mellitus, such as IDDM, NIDDM, gestational diabetes, juvenile diabetes, and the like, using combination therapy. In certain aspects, the pharmaceutical compositions comprise a pharmaceutically acceptable carrier with a compound of Formula I and an antidiabetic agent. Advantageously, the compositions of the present invention provide clinical advantage over the use of a single agent alone. As such, the present invention provides a composition comprising:

i) a compound of Formula I



wherein

Ar^1 is an aryl group; X is a divalent linkage selected from $(\text{C}_1\text{-C}_6)\text{alkylene}$, $(\text{C}_1\text{-C}_6)\text{alkylenoxy}$, $(\text{C}_1\text{-C}_6)\text{alkylenamino}$, $(\text{C}_1\text{-C}_6)\text{alkylene-S(O)}_k\text{-}$, -O- , -C(O)- , $\text{-N(R}^{11}\text{)-}$, $\text{-N(R}^{11}\text{)C(O)-}$, $\text{-S(O)}_k\text{-}$ and a single bond, wherein R^{11} is a member selected from hydrogen, $(\text{C}_1\text{-C}_8)\text{alkyl}$, $(\text{C}_1\text{-C}_8)\text{heteroalkyl}$ and $\text{aryl(C}_1\text{-C}_4\text{)alkyl}$; and the

subscript k is an integer of from 0 to 2; Y is a divalent linkage selected from alkylene, -O-, -C(O)-, -N(R¹²)-S(O)_m-, -N(R¹²)-S(O)_m-N(R¹³)-, -N(R¹²)C(O)-, -S(O)_n- and a single bond, wherein R¹² and R¹³ are members independently selected from hydrogen, (C₁-C₈)alkyl, (C₁-C₈)heteroalkyl and aryl(C₁-C₄)alkyl; and the subscripts m and n are
 5 independently integers of from 0 to 2;

R¹ is a member selected from the group hydrogen, heteroalkyl, aryl, arylalkyl, halogen, cyano, nitro, (C₁-C₈)alkyl, (C₁-C₈)alkoxy, -C(O)R¹⁴, -CO₂R¹⁴, C(O)NR¹⁵R¹⁶, -S(O)_p-R¹⁴, -S(O)_q-NR¹⁵R¹⁶, -O-C(O)-OR¹⁷, -O-C(O)-R¹⁷, -O-C(O)-NR¹⁵R¹⁶, -N(R¹⁴)-C(O)-NR¹⁵R¹⁶, -N(R¹⁴)-C(O)-R¹⁷ and -N(R¹⁴)-C(O)-OR¹⁷; wherein
 10 R¹⁴ is a member selected from hydrogen, (C₁-C₈)alkyl, (C₁-C₈)heteroalkyl, aryl and aryl(C₁-C₄)alkyl;

R¹⁵ and R¹⁶ are members independently selected from hydrogen, (C₁-C₈)alkyl, (C₁-C₈)heteroalkyl, aryl, and aryl(C₁-C₄)alkyl, or taken together with the nitrogen to which each is attached form a 5-, 6- or 7-membered ring;

15 R¹⁷ is a member selected from alkyl, heteroalkyl, aryl and arylalkyl;

the subscript p is an integer of from 0 to 3;

the subscript q is an integer of from 1 to 2;

R² is a member selected from (C₁-C₈)alkyl, (C₁-C₈)heteroalkyl, aryl and aryl(C₁-C₄)alkyl;

20 R³ is a member selected from hydrogen, halogen, cyano, nitro, (C₁-C₈)alkyl and (C₁-C₈)alkoxy;

including pharmaceutically acceptable salts of compounds of Formula I;
 and

ii) one or more antidiabetic agents, including, but not limited to,
 25 sulfonylureas, biguanides, glitazones and other PPAR_γ agonists, PPAR_α agonists, PPAR_δ agonists, α-glucosidase inhibitors, potassium channel antagonists, aldose reductase inhibitors, glucagon antagonists, activators of RXR, insulin therapy or other anti-obesity agent (5), prodrugs thereof, or pharmaceutically acceptable salts of the antidiabetic agents, and a pharmaceutically acceptable carrier or diluent.

30 In certain aspects, the compositions of the present invention comprise a compound of Formula I formulated together with one or more antidiabetic agents. Alternatively, the compositions of the present invention comprise a compound of Formula

I independently formulated with one or more antidiabetic agents *i.e.*, separately formulated.

Suitable antidiabetic agents include, but are not limited to, sulfonylureas, biguanides, glitazones and other PPAR γ agonists, α -glucosidase inhibitors, potassium channel antagonists, aldose reductase inhibitors, glucagon antagonists, activators of RXR, 5 activators of PPAR α , activators of PPAR δ , insulin therapy or other anti-obesity agents. The administration of a composition comprising i) a compound of Formula I, which are PPAR γ modulators and known to increase peripheral tissue sensitivity to insulin, with ii) an antidiabetic agent such as insulin therapy, or a stimulator of insulin secretion, and the 10 like, increases the efficacy of either agent alone. In addition to increased efficacy, the combination therapy of the present invention allows for a concomitant reduction in the dose of the agents. The combination therapy of a compound of Formula I and one or more of another antidiabetic agents (*e.g.*, biguanides, glitazones, RXR ligands, PPAR γ agonists, *etc.*) results in a reduction in the side effects normally associated with certain 15 antidiabetic agents.

In certain aspects, compounds of Formula I are administered in combination with antidiabetic agents that are ineffective for stimulation of insulin secretion or insulin sensitivity, such as α -glucosidase inhibitors, potassium channel antagonists, aldose reductase inhibitors, glucagon antagonists, RXR ligands, PPAR α 20 agonists, PPAR δ agonists, and anti-obesity agents. Surprisingly, these types of combination therapy result in enhanced efficacy relative to the use of the single agents alone.

In another embodiment, the present invention provides methods of treating metabolic or inflammatory disorders in a host by administering a composition of the 25 present invention. In certain preferred aspects, the method includes the administration of a composition comprising a combination of a compound of Formula I with the antidiabetic agent delivered in a simultaneous manner, such as in a single formulation. In certain other aspects, the methods of the present invention include combination therapy wherein the compound of Formula I is administered first in one formulation, followed by 30 the antidiabetic agent in a separate formulation. The methods also include an antidiabetic agent being delivered first in one formulation, followed by a compound of Formula I in a separate formulation. The present invention includes all such methods of administration. The combination therapy is especially efficacious on conditions associated with diabetes,

such as obesity, cardiovascular disease, hypercholesterolemia and other lipid disorders, peripheral neuropathies and other neurological disorders, and the like.

These and other feature and advantages will become more apparent when read with the accompanying Figures and detailed description that follow.

5

DEFINITIONS

The term "alkyl," by itself or as part of another substituent, means, unless otherwise stated, a straight or branched chain, or cyclic hydrocarbon radical, or combination thereof, which may be fully saturated, mono- or polyunsaturated and can include di- and multivalent radicals, having the number of carbon atoms designated (*i.e.* C₁-C₁₀ means one to ten carbons). Examples of saturated hydrocarbon radicals include groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, isobutyl, sec-butyl, cyclohexyl, (cyclohexyl)ethyl, cyclopropylmethyl, homologs and isomers of, for example, n-pentyl, n-hexyl, n-heptyl, n-octyl, and the like. An unsaturated alkyl group is one having one or more double bonds or triple bonds. Examples of unsaturated alkyl groups include vinyl, 2-propenyl, crotyl, 2-isopentenyl, 2-(butadienyl), 2,4-pentadienyl, 3-(1,4-pentadienyl), ethynyl, 1- and 3-propynyl, 3-butylnyl, and the higher homologs and isomers. The term "alkyl," unless otherwise noted, is also meant to include those derivatives of alkyl defined in more detail below as "heteroalkyl," "cycloalkyl" and "alkylene." The term "alkylene" by itself or as part of another substituent means a divalent radical derived from an alkane, as exemplified by -CH₂CH₂CH₂CH₂-. Typically, an alkyl group will have from 1 to 24 carbon atoms, with those groups having 10 or fewer carbon atoms being preferred in the present invention. A "lower alkyl" or "lower alkylene" is a shorter chain alkyl or alkylene group, generally having eight or fewer carbon atoms. An alkanoyl is a RCO- group.

The term "heteroalkyl," by itself or in combination with another term, means, unless otherwise stated, a stable straight or branched chain, or cyclic hydrocarbon radical, or combinations thereof, consisting of the stated number of carbon atoms and from one to three heteroatoms selected from the group consisting of O, N, Si and S, and wherein the nitrogen and sulfur atoms can optionally be oxidized and the nitrogen heteroatom can optionally be quaternized. The heteroatom(s) O, N and S may be placed at any interior position of the heteroalkyl group. The heteroatom Si may be placed at any position of the heteroalkyl group, including the position at which the alkyl group is attached to the remainder of the molecule. Examples include -CH₂-CH₂-O-CH₃, -CH₂-

CH₂-NH-CH₃, -CH₂-CH₂-N(CH₃)-CH₃, -CH₂-S-CH₂-CH₃, -CH₂-CH₂-S(O)-CH₃, -CH₂-CH₂-S(O)₂-CH₃, -CH=CH-O-CH₃, -Si(CH₃)₃, -CH₂-CH=N-OCH₃, and -CH=CH-N(CH₃)-CH₃. Up to two heteroatoms may be consecutive, such as, for example, -CH₂-NH-OCH₃ and -CH₂-O-Si(CH₃)₃. Also included in the term "heteroalkyl" are those radicals

5 described in more detail below as "heteroalkylene" and "heterocycloalkyl." The term "heteroalkylene" by itself or as part of another substituent means a divalent radical derived from heteroalkyl, as exemplified by -CH₂-CH₂-S-CH₂CH₂- and -CH₂-S-CH₂-CH₂-NH-CH₂-. For heteroalkylene groups, heteroatoms can also occupy either or both of the chain termini. Still further, for alkylene and heteroalkylene linking groups, no orientation
10 of the linking group is implied.

The terms "cycloalkyl" and "heterocycloalkyl", by themselves or in combination with other terms, represent, unless otherwise stated, cyclic versions of "alkyl" and "heteroalkyl", respectively. Additionally, for heterocycloalkyl, a heteroatom can occupy the position at which the heterocycle is attached to the remainder of the
15 molecule. Examples of cycloalkyl include cyclopentyl, cyclohexyl, 1-cyclohexenyl, 3-cyclohexenyl, cycloheptyl, and the like. Examples of heterocycloalkyl include 1 - (1,2,5,6-tetrahydropyridyl), 1 - piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-morpholinyl, 3-morpholinyl, tetrahydrofuran-2-yl, tetrahydrofuran-3-yl, tetrahydrothien-2-yl, tetrahydrothien-3-yl, 1-piperazinyl, 2-piperazinyl, and the like.

20 The terms "halo" or "halogen," by themselves or as part of another substituent, mean, unless otherwise stated, a fluorine, chlorine, bromine, or iodine atom. Additionally, terms such as "fluoroalkyl," are meant to include monofluoroalkyl and polyfluoroalkyl.

The term "aryl," employed alone or in combination with other terms (*e.g.*,
25 aryloxy, arylthioxy, arylalkyl, aroyl (ArCO), heteroaroyl) means, unless otherwise stated, an aromatic substituent which can be a single ring or multiple rings (up to three rings) which are fused together or linked covalently. The rings may each contain from zero to four heteroatoms selected from N, O, and S, wherein the nitrogen and sulfur atoms are optionally oxidized, and the nitrogen atom(s) are optionally quaternized. The aryl groups
30 that contain heteroatoms may be referred to as "heteroaryl" and can be attached to the remainder of the molecule through a heteroatom. Non-limiting examples of aryl groups include aminobenzoheteroazolyl, 2-azanaphthalenyl, bezoxazolyl, phenyl, 1-naphthyl, 2-naphthyl, 4-biphenyl, 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 3-pyrazolyl, 2-imidazolyl, 4-imidazolyl, pyrazinyl, 2-oxazolyl, 4-oxazolyl, 2-phenyl-4-oxazolyl, 5-oxazolyl, 3-

isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidyl, 4-pyrimidyl, 5-benzothiazolyl, purinyl, 2-benzimidazolyl, 5-indolyl, 1-isoquinolyl, 5-isoquinolyl, 2-quinoxaliny, 5-quinoxaliny, 3-quinolyl, 6-quinolyl, thiobenzoxazolyl, thiobenzothiazolyl and thiobenzimidazolyl. Substituents for each of the above noted aryl ring systems are selected from the group of acceptable substituents described below. The term "arylalkyl" is meant to include those radicals in which an aryl group is attached to an alkyl group (*e.g.*, benzyl, phenethyl, pyridylmethyl and the like) or a heteroalkyl group (*e.g.*, phenoxymethyl, 2-pyridyloxymethyl, 3-(1-naphthyloxy)propyl, and the like).

Each of the above terms (*e.g.*, "alkyl," "heteroalkyl" and "aryl") are meant to include both substituted and unsubstituted forms of the indicated radical. Preferred substituents for each type of radical are provided below.

Substituents for the alkyl and heteroalkyl radicals (including those groups often referred to as alkylene, alkenyl, heteroalkylene, heteroalkenyl, alkynyl, cycloalkyl, heterocycloalkyl, cycloalkenyl, and heterocycloalkenyl) can be a variety of groups selected from: -OR', =O, =NR', =N-OR', -NR'R'', -SR', -halogen, -SiR'R''R''', -OC(O)R', -C(O)R', -CO₂R', CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -NR'-C(O)NR''R''', -NR''C(O)₂R', -NH-C(NH₂)=NH, -NR'C(NH₂)=NH, -NH-C(NH₂)=NR', -S(O)R', S(O)₂R', -S(O)₂NR'R'', -CN and -NO₂ in a number ranging from zero to (2N+1), where N is the total number of carbon atoms in such radical. R', R'' and R''' each independently refer to hydrogen, unsubstituted(C₁-C₈)alkyl and heteroalkyl, unsubstituted aryl, aryl substituted with 1-3 halogens, unsubstituted alkyl, alkoxy or thioalkoxy groups, or aryl-(C₁-C₄)alkyl groups. When R' and R'' are attached to the same nitrogen atom, they can be combined with the nitrogen atom to form a 5-, 6-, or 7-membered ring. For example, -NR'R'' is meant to include 1-pyrrolidinyl and 4-morpholinyl. From the above discussion of substituents, one of skill in the art will understand that the term "alkyl" is meant to include groups such as haloalkyl (*e.g.*, -CF₃ and -CH₂CF₃) and acyl (*e.g.*, -C(O)CH₃, -C(O)CF₃, -C(O)CH₂OCH₃, and the like).

Similarly, substituents for the aryl groups are varied and are selected from: --halogen, -OR', -OC(O)R', -NR'R'', -SR', -R', -CN, -NO₂, -CO₂R', -CONR'R'', -C(O)R', -OC(O)NR'R'', -NR''C(O)R', -NR''C(O)₂R', -NR'-C(O)NR''R''', -NH-C(NH₂)=NH, -NR'C(NH₂)=NH, -NH-C(NH₂)=NR', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -N₃, -CH(Ph)₂, perfluoro(C₁-C₄)alkoxy, and perfluoro(C₁-C₄)alkyl, in a number ranging from zero to the total number of open valences on the aromatic ring system; and where

R', R'' and R''' are independently selected from hydrogen, (C₁-C₈)alkyl and heteroalkyl, unsubstituted aryl, (unsubstituted aryl)-(C₁-C₄)alkyl, and (unsubstituted aryl)oxy-(C₁-C₄)alkyl.

Two of the substituents on adjacent atoms of the aryl ring may optionally be replaced with a substituent of the formula -T-C(O)-(CH₂)_q-U- wherein T and U are independently -NH-, -O-, -CH₂- or a single bond, and q is an integer of from 0 to 2. Alternatively, two of the substituents on adjacent atoms of the aryl ring may optionally be replaced with a substituent of the formula -A-(CH₂)_r-B-, wherein A and B are independently -CH₂-, -O-, -NH-, -S-, -S(O)-, -S(O)₂-, -S(O)₂NR'- or a single bond, and r is an integer of from 1 to 3. One of the single bonds of the new ring so formed may optionally be replaced with a double bond. Alternatively, two of the substituents on adjacent atoms of the aryl ring may optionally be replaced with a substituent of the formula -(CH₂)_s-X-(CH₂)_t-, where s and t are independently integers of from 0 to 3, and X is -O-, -NR'-, -S-, -S(O)-, -S(O)₂-, or -S(O)₂NR'-. The substituent R' in -NR'- and -S(O)₂NR'- is selected from hydrogen or unsubstituted (C₁-C₆)alkyl.

In certain instances, aryl means two aryl groups joined by a heteroatom. These groups include diphenyl ether, phenoxy substituted 2-azanaphthalene, phenyl-thiobenzothiazole, phenyl-heterobenzoxazoles, phenyl-thiobenzoimidazoles and phenyl-heterobenzoheteroazoles (see structures IIa-IIf below).

As used herein, the term "heteroatom" is meant to include oxygen (O), nitrogen (N), sulfur (S) and silicon (Si).

The term "pharmaceutically acceptable salts" is meant to include salts of the active compounds which are prepared with relatively nontoxic acids or bases, depending on the particular substituents found on the compounds described herein. When compounds of the present invention contain relatively acidic functionalities, base addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired base, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable base addition salts include sodium, potassium, calcium, ammonium, organic amino, or magnesium salt, or a similar salt. When compounds of the present invention contain relatively basic functionalities, acid addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired acid, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable acid addition salts include, but are not limited to, those derived from inorganic acids like hydrochloric, hydrobromic, nitric, carbonic, monohydrogencarbonic,

phosphoric, monohydrogenphosphoric, dihydrogenphosphoric, sulfuric, monohydrogensulfuric, hydriodic, or phosphorous acids and the like, as well as the salts derived from relatively nontoxic organic acids like acetic, propionic, isobutyric, oxalic, maleic, malonic, benzoic, succinic, suberic, fumaric, mandelic, phthalic, benzenesulfonic, p-tolylsulfonic, citric, tartaric, methanesulfonic, and the like. Also included are salts of amino acids such as arginate and the like, and salts of organic acids like glucuronic or galactunoric acids and the like (*see, for example, Berge, S.M., et al., "Pharmaceutical Salts", Journal of Pharmaceutical Science, 1977, 66, 1-19*). Certain specific compounds of the present invention contain both basic and acidic functionalities that allow the compounds to be converted into either base or acid addition salts.

The neutral forms of the compounds can be regenerated by contacting the salt with a base or acid and isolating the parent compound in the conventional manner. The parent form of the compound differs from the various salt forms in certain physical properties, such as solubility in polar solvents, but otherwise the salts are equivalent to the parent form of the compound for the purposes of the present invention.

In addition to salt forms, the present invention provides compounds that are in a prodrug form. Prodrugs of the compounds described herein are those compounds that readily undergo chemical changes under physiological conditions to provide the compounds of the present invention. Additionally, prodrugs can be converted to the compounds of the present invention by chemical or biochemical methods in an *ex vivo* environment. For example, prodrugs can be slowly converted to the compounds of the present invention when placed in a transdermal patch reservoir with a suitable enzyme or chemical reagent.

Certain compounds of the present invention can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms are equivalent to unsolvated forms and are intended to be encompassed within the scope of the present invention. Certain compounds of the present invention can exist in multiple crystalline or amorphous forms. In general, all physical forms are equivalent for the uses contemplated by the present invention and are intended to be within the scope of the present invention.

Certain compounds of the present invention possess asymmetric carbon atoms (optical centers) or double bonds; the racemates, diastereomers, geometric isomers and individual isomers are all intended to be encompassed within the scope of the present invention.

The compounds of the present invention can also contain unnatural proportions of atomic isotopes, stable isotopes *etc.*, at one or more of the atoms that constitute such compounds. For example, the compounds may be radiolabeled with radioactive isotopes, such as for example tritium (^3H), iodine-125 (^{125}I) or carbon-14 (^{14}C).

5 All isotopic variations of the compounds of the present invention, whether radioactive or not, are intended to be encompassed within the scope of the present invention.

The term "prodrug" refers to compounds that are drug precursors, which, following administration, release the drug *in vivo* via a chemical or physiological process (e.g., a prodrug on being brought to the physiological pH is converted to the desired drug
10 form).

"A combination amount sufficient," "an effective combination amount" "therapeutically effective combination amount" or "an effective amount of the combination of" all refer to a combined amount of both a compound of Formula I and the antidiabetic agent that is effective to ameliorate symptoms associated with diabetic
15 diseases. As used herein, the term "combination" of compound of Formula I with an antidiabetic agent means the two compounds can be delivered in a simultaneous manner, in combination therapy wherein the compound of Formula I is administered first, followed by the antidiabetic agent, as well as wherein the antidiabetic agent is delivered first, followed by a compound of Formula I. The desired result can be either a subjective
20 relief of a symptom(s) or an objectively identifiable improvement in the recipient of the dosage.

The terms "synergistic effective amount" refers to a combined amount of both a compound of Formula I and an antidiabetic agent that is effective to cause a synergistic effect. Synergy is a biological phenomenon in which the effectiveness of two
25 active components in a mixture is more than additive, *i.e.*, the effectiveness is greater than the equivalent concentration of either component alone. In certain aspects, the effectiveness of the combination therapy of a compound of Formula I and an antidiabetic agent is synergistic. Thus, synergism is a result, or function, that is more than the sum of the results, or functions of individual elements.

30 The term "simultaneous manner" and "combination treatment" refer to an administration protocol wherein the compound of the present invention and at least one antidiabetic agent are administered within a single 24-hour period.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows an illustrative scheme for synthesizing compounds of Formula I.

Figure 2 shows illustrative antidiabetic agents.


Figure 3 illustrates various compounds of Formula I.

DETAILED DESCRIPTION OF THE INVENTION AND PREFERRED EMBODIMENTS

COMPOSITIONS

In one embodiment, the present invention provides a pharmaceutical composition comprising (i) a compound of Formula I and (ii) an antidiabetic agent. Advantageously, the compositions of the present invention provide clinical advantage over the use of a single agent alone. As such, the present invention provides a composition comprising:

i) a compound of Formula I



In Formula I, Ar^1 is a functional group including, but not limited to, an aryl group. X , in Formula I, is a divalent linkage including, but not limited to, (C_1-C_6) alkylene, (C_1-C_6) alkylenoxy, (C_1-C_6) alkylenamino, (C_1-C_6) alkylene- $S(O)_k$ -, $-O$ -, $-C(O)$ -, $-N(R^{11})$ -, $-N(R^{11})C(O)$ -, $-S(O)_k$ - and a single bond. R^{11} , in Formula I, is a functional group including, but not limited to, hydrogen, (C_1-C_8) alkyl, (C_1-C_8) heteroalkyl and aryl(C_1-C_4)alkyl. The index "k" is an integer of from 0 to 2 inclusive.

Y , in Formula I, is a divalent linkage including, but not limited to, alkylene, $-O$ -, $-C(O)$ -, $-N(R^{12})-S(O)_m$ -, $-N(R^{12})-S(O)_m-N(R^{13})$ -, $-N(R^{12})C(O)$ -, $-S(O)_n$ - and a single bond. R^{12} and R^{13} , in Formula I, are each independent functional groups including, but not limited to, hydrogen, (C_1-C_8) alkyl, (C_1-C_8) heteroalkyl and aryl(C_1-C_4)alkyl. The indices "m" and "n" are each independent integers of from 0 to 2 inclusive.

In Formula I, R^1 , is a functional group including, but not limited to, hydrogen, heteroalkyl, aryl, arylalkyl, halogen, cyano, nitro, (C_1-C_8) alkyl, (C_1-C_8) alkoxy,

-C(O)R¹⁴, -CO₂R¹⁴ and -C(O)NR¹⁵R¹⁶, -S(O)_p-R¹⁴, -S(O)_q-NR¹⁵R¹⁶, -O-C(O)-OR¹⁷, -O-C(O)-R¹⁷, -O-C(O)-NR¹⁵R¹⁶, -N(R¹⁴)-C(O)-NR¹⁵R¹⁶, -N(R¹⁴)-C(O)-R¹⁷ and -N(R¹⁴)-C(O)-OR¹⁷. R¹⁴, in Formula I, is a functional group including, but not limited to, hydrogen, (C₁-C₈)alkyl, (C₁-C₈)heteroalkyl, aryl and aryl(C₁-C₄)alkyl. R¹⁵ and R¹⁶, in
 5 Formula I, are each independent functional groups including, but not limited to, hydrogen, (C₁-C₈)alkyl, (C₁-C₈)heteroalkyl, aryl, and aryl(C₁-C₄)alkyl. In an alternative embodiment, R¹⁵ and R¹⁶ taken together with the nitrogen to which each is bound form a 5-, 6- or 7-membered ring. In Formula I, R¹⁷, is a functional group including, but not limited to, alkyl, heteroalkyl, aryl and arylalkyl. The index "p" is an integer of from 0 to
 10 3 inclusive. The index "q" is an integer of from 1 to 2.

In Formula I, R², is a functional group including, but not limited to, (C₁-C₈)alkyl, heteroalkyl, (C₁-C₈)heteroalkyl, aryl and aryl(C₁-C₄)alkyl. R³, in Formula I, is a functional group including, but not limited to hydrogen, a halogen, cyano, nitro and (C₁-C₈)alkoxy.

15 Based on the unique features of the compounds of Formula I, the combination of one of these compounds with one or more antidiabetic agents described herein provides a significant clinical advantage over the use of a single agent alone. Thus, (1) the combination of a compound of Formula I (which as activator of PPAR_γ, is thought to increase peripheral tissue sensitivity to insulin) with either insulin therapy, or a
 20 stimulator of insulin secretion (*e.g.*, a sulfonylurea) increases the efficacy of either agent alone, and moreover, allows for the reduction in dosage of all agents used in the combination therapy. In addition, (2) the combination therapy between a compound of Formula I and one or more other agents that increase insulin sensitivity (*e.g.*, biguanides, glitazones, RXR ligands, PPAR_γ agonists, and the like.), results in an enhanced effect
 25 between the various agents, with reduction in the side effects normally associated with these other agents. Further, (3) compounds of Formula I can be administered in combination with antidiabetic agents whose mode of action is other than stimulation of insulin secretion or insulin sensitivity (*e.g.*, α-glucosidase inhibitors, potassium channel antagonists, aldose reductase inhibitors, glucagon antagonists, RXR ligands and anti-obesity agents). Importantly, these types of combinations result in enhanced efficacy
 30 relative to the use of a single agent alone. In addition, the present invention includes (4) a combination treatment comprising a compound of Formula I in combination with agents aimed at treating any one of the conditions often associated with diabetes, such as obesity,

cardiovascular disease, hypercholesterolemia and other lipid disorders, peripheral neuropathies and other neurological disorders.

The compounds of Formula I are known to interact with the peroxisome proliferator-activated receptor gamma (PPAR γ) protein, a member of the nuclear receptor family. These compounds possess the ability to bind to PPAR γ with high affinity while inducing only a partial functional activation of the receptor's potential transcriptional activity. This property is sometimes described as "partial agonism". The exact potency and intrinsic agonist activity of each compound of Formula I is a function of the compound's structure in a relatively predictable manner.

Compounds of Formula I, as partial PPAR γ agonists, have been shown, in diabetic animal models, to possess the anti-diabetic efficacy of the glitazones, but with a decreased propensity to induce body weight gain. One of the major side effects of glitazones (*e.g.*, troglitazone, rosiglitazone and pioglitazone) is their obesity-inducing potential. This effect is thought to be related to their full activation of PPAR γ receptors in adipocytes and pre-adipocytes, and it is not thought to be required for anti-diabetic efficacy. Advantageously, compounds of Formula I possess a diminished propensity towards inducing body weight gain and thus, one of the major side effects of known glitazones. Moreover, compounds of Formula I have also shown beneficial effects on plasma lipid profiles.

In one embodiment, compounds of Formula I are described in U.S. Provisional Patent Application Serial Number 60/141,672, filed June 30, 1999; WO 01/00579, published January 4, 2001; to McGee *et al.*, and incorporated herein by reference in their entirety for all purposes. As described therein, compounds of Formula I can be synthesized by a variety of methods, some of which are illustrated in Figure 1. With reference to Figure 1, using commercial available 2-chloro-5-nitrobenzonitrile (i) and treatment of (i) with a phenol, thiophenol, or optionally protected aniline in the presence of base and heat generates adduct (ii). Reduction of the nitro group in (ii) with, for example, hydrogen in the presence of Raney nickel catalyst provides an aniline derivative (iii). Sulfonylation of (iii) with an appropriate arylsulfonyl halide in the presence of base (typically a tertiary amine) provides target compound (iv). Compound (iii) can also be converted to a related compound of formula (vi) in which the orientation of the sulfonamide linkage is reversed. Thus, conversion of the aniline (iii) to the benzenesulfonyl chloride (v) can be accomplished using methods described in Hoffman, *Organic Syntheses Collective Volume VII*, p. 508-511. Subsequent treatment of (v) with




an appropriate aniline provides the target compound (vi). Other compounds of the present invention can be prepared beginning with, for example, 3,4-difluoronitrobenzene, 3-chloro-4-fluoronitrobenzene, 2-chloro-5-nitroanisole, 3-bromo-4-fluoronitrobenzene and the like.

Moreover, in certain embodiments, compounds of Formula I are described in WO 99/38845, published August 5, 1999, to De La Brouse-Elwood *et al.*, and incorporated herein by reference in its entirety for all purposes. Certain compounds as described therein, are potent, selective activators of PPAR γ and are useful for the treatment of NIDDM and other disorders related to lipid metabolism and energy homeostasis. In other aspects, and depending on the biological environment (*e.g.*, cell type, pathological condition of the host, and the like.), the compounds can activate or block the actions of PPAR γ . By activating the PPAR γ receptor, the compounds will find use as therapeutic agents capable of modulating conditions mediated by the PPAR γ receptor. Additionally, the compounds are useful for the prevention and treatment of complications of diabetes (*e.g.*, neuropathy, retinopathy, glomerulosclerosis, and cardiovascular disorders), and treating hyperlipidemia. The compounds are useful for the modulation of inflammatory conditions which most recently have been found to be controlled by PPAR γ (*see*, Ricote, *et al.*, *Nature*, 391:79-82 (1998) and Jiang, *et al.*, *Nature*, 391:82-86 (1998)). Examples of inflammatory conditions include rheumatoid arthritis and atherosclerosis. Compounds that act via antagonism of PPAR γ are useful for treating obesity, hypertension, hyperlipidemia, hypercholesterolemia, hyperlipoproteinemia, and metabolic disorders.

Preferred compounds of Formula I are those in which Ar¹ is a pyridyl ring having a single substituent selected from the group consisting of halogen, -OCF₃ and -CF₃; X is a divalent linkage selected from the group of -O-, -SO₂-, -S-, -C(O)-, -CH₂- and combinations thereof; Y is a divalent linkage selected from the group of -NH-S(O)₂- and -NH-C(O)-; R¹ is selected from the group consisting of hydrogen, halogen, cyano, (C₁-C₈)alkyl, (C₁-C₈)alkoxy, -C(O)(C₁-C₈ alkyl) and -C(O)NR¹⁵R¹⁶ in which R¹⁵ and R¹⁶ are selected from hydrogen, (C₁-C₈)alkyl, aryl and aryl(C₁-C₄)alkyl; R² is a phenyl or pyridyl ring, optionally substituted by 0-3 groups selected from halogen, (C₁-C₈)alkyl, -O-(C₁-C₈)alkyl, CF₃, and -CN; and R³ is hydrogen, halogen, cyano or (C₁-C₄)alkoxy.

One of skill in the art will understand that a number of structural isomers are represented by Formula I. Preferred isomers are those in which the groups on the

phenyl ring occupy positions that are not all contiguous. Particularly preferred compounds are those having the structural orientations represented by the formulae:




5


Still further preferred are those compounds having the structural orientation represented by formula Ia or Ib. Still other preferred compounds, are those of formula Ia or Ib in which the positions of R¹ and R³ are switched (or reversed).

Yet other preferred compounds are those in which Ar¹-X- and -Y-R² occupy positions para to one another (exemplified by Ii).


10



Still another group of preferred compounds are represented by the formula:



Equally preferred Ar¹ moieties are set forth below. Each of the rings below can be optionally substituted with one or more functional groups. In the structures below, "W" is a heteroatom such as N, O or S.



The compositions of the present invention further comprises one or more (ii) antidiabetic compounds. A wide range of antidiabetic agents can be used in the compositions and methods of the present invention. Suitable agents include, but are not limited to, one or more antidiabetic agent such as sulfonylureas, biguanides, glitazones and other PPAR γ agonists, α -glucosidase inhibitors, potassium channel antagonists, aldose reductase inhibitors, glucagon antagonists, activators of RXR, insulin therapy or other anti-obesity agent, prodrugs thereof, or pharmaceutically acceptable salts of the antidiabetic agents. In certain instances, the antidiabetic agents include, but are not limited to, one or more agents such as insulin and insulin analogs (*e.g.* LysPro insulin); GLP-1 (7-37) (insulinotropin) and GLP-1 (7-36)-NH₂; sulfonylureas and analogs, including, but not limited to, chlorpropamide, glibenclamide, tolbutamide, tolazamide, acetohexamide, glipizide, glimepiride, repaglinide and meglitinide; biguanides, including, but not limited to, metformin, phenformin and buformin.


In another embodiment, the antidiabetic agents include various forms of insulin, such as insulin in its various dosage forms, subcutaneous, oral, inhaled, and the like, molecular variations, and short-, medium- and long-acting versions. Suitable insulin sources include, but are not limited to, recombinant human insulin, natural pig insulin,

natural ox insulin, natural bovine insulin, natural human insulin, recombinant human argine-insulin, recombinant human aspartic-insulin, dalanated insulin, defalan insulin, glargine insulin, human insulin zinc, human insulin isophane, lispro insulin, neutral insulin, human proinsulin, and the like.

5 In one embodiment, the antidiabetic agents used in the compositions of the present invention are sulfonylureas. Preferred sulfonylureas suitable for use in the present invention include, but are not limited to, acetohexamide, chlorpropamide, glyburide, glipizide, gliclazide, glimepiride, gliquidone, glisoxepid, glibornuride, gliamilide, glibornuride, glicetanile, gliflumide, glymidine, glyparamide, tolpyrramide, 10 glyhexamide, phenbutamide, tolazamide, tolbutamide and tolcyclamide. Those of skill in the art will know of other sulfonylureas suitable for use in the present invention.

Sulfonylureas are a group of drugs that induce hypoglycemia by stimulating insulin release from the pancreas. Generally, sulfonylureas have found wide utility in the treatment of NIDDM. Their efficacy is decreased in IDDM because of the 15 inherent inability of the patient to produce insulin. Adverse reactions to sulfonylureas occur in a fraction of patients, particularly the elderly. One of the most severe side effects is hypoglycemia and coma. Other side effects include nausea and vomiting, cholestatic jaundice, agranulocytosis, cardiovascular mortality, aplastic and hemolytic anemias, generalized hypersensitivity reactions and dermatological reactions. The compounds of 20 Formula I increase insulin sensitivity, and when used in combination with sulfonylureas, increase the efficacy and side effect profile of either type of agent alone. In certain instances, the combination therapy allows the use of lower doses of both agents.

In certain embodiments, the sulfonylurea used in the compositions of the present invention have the following formula:



sulfonylurea structure

II


25

wherein R is H, halogen, alkyl, alkanoyl, aryl, arylalkyl, heteroaryl, aroylaminoalkyl, heteroaroylaminoalkyl, or cycloalkyl; and R' is alkyl, cycloalkyl and heterocycloalkyl (including those that are listed to the rest of the molecule via a heteroatom that is part of R'). Preferred sulfonylureas are those in which R is 4-chloro, 4-methyl, 4-acetyl, 4-[2- 30 [(3-chloro-5-methoxybenzoyl)amino]ethyl] or 4-[2-[(5-methyl-2-

pyrazinylcarbonyl)amino]ethyl]; and R' is propyl, butyl, 1-piperidyl, cyclohexyl, cycloheptyl, 4-dimethylaminophenyl, 1-hexahydroazepine, or 3-azabicyclo[3.3.0]oct-3-yl.

In another embodiment, the antidiabetic agents used in the compositions of the present invention are biguanides. Biguanides are a group of drugs that are efficacious in the treatment of hyperglycemia. Preferred biguanides suitable for use in the present invention include, but are not limited to, metformin, phenformin and buformin. Unlike the sulfonylureas, metformin does not induce release of insulin from the pancreas. Without being bound by any particular theory, it is thought that its effects are mediated by increasing insulin activity in peripheral tissues, reducing hepatic glucose output due to inhibition of gluconeogenesis and reducing the absorption of glucose from the intestine. Side effects associated with the use of biguanides include lactic acidosis, diarrhea, nausea, and anorexia.

In certain aspects, the biguanides used in the compositions of the present invention have the following formula:




biguanides

III

wherein R, R' and R'' are each selected independently from H, alkyl, cycloalkyl, aryl, arylalkyl heteroalkyl, heteroaryl, heteroarylalkyl and aryl-heteroalkyl. Preferably, R, R' and R'' are each independently selected from hydrogen, (C₁-C₃)alkyl and aryl-(C₁-C₃)alkyl. Preferred biguanides include metformin, buformin, etoformin and phenformin.

In another embodiment, the antidiabetic agent is a glitazone. Glitazones, also known as thiazolidinediones, produce antidiabetic effects by modulating the activity of the nuclear receptor PPAR γ . Particularly preferred members of this group of antidiabetic agents have the following formula:



glitazone structure

IV

wherein R is a functional group containing from six to twenty carbon atoms and at least one aromatic or non-aromatic ring system; and X and Y are each selected independently from O, S or NH. Suitable glitazone compounds are disclosed in *Current Pharmaceutical Design*, 1996; 2:85-101, incorporated herein by reference, and include compounds such as troglitazone, rosiglitazone, pioglitazone, JTT-501, YM-440, ciglitazone, darglitazone, englitazone, AD-5075, and BM-131246. Preferably, the glitazone compounds used in the present invention include troglitazone, rosiglitazone, and pioglitazone. Other PPAR γ activators that are not part of the glitazone family, such as GW-409544 and GI-262570, can be used. Figure 2 sets forth PPAR γ modulators suitable for use in the present invention.

In certain aspects, PPAR γ modulators are disclosed in WO 99/58510, published November 18, 1999, to Momose *et al.*, and incorporated herein by reference in its entirety for all purposes. As disclosed therein, oxyiminoalkanoic acid derivatives are disclosed that are useful for treating and/or preventing diabetes mellitus, hyperlipemia, impaired glucose tolerance, inflammatory disease and arteriosclerosis. Moreover, the compounds are useful for controlling retinoid-related receptors, enhancing insulin sensitivity and improving insulin resistance, and for treating and preventing diabetic complications. Other PPAR γ modulators are disclosed in WO 99/38850, published August 5, 1999, to Lohray *et al.*, and incorporated herein by reference in its entirety for all purposes.

In another embodiment, the present invention provides compositions comprising compounds of Formula I and α -glucosidase inhibitors. Preferred α -glucosidase inhibitors include acarbose, celgosivir, camigliabase, voglibose and miglitol. Suitable α -glucosidase inhibitors are disclosed in WO 99/29327, to Odaka *et al.*, published June 17, 1999, entitled "Use of an α -glucosidase inhibitor to treat high-risk impaired glucose tolerance" and incorporated herein by reference in its entirety for all purposes. As described therein, the α -glucosidase inhibitor is particularly useful in the treatment of non-insulin-dependent diabetes mellitus. In addition, WO 99/26606, published June 3, 1999, to Goldman *et al.*, and entitled "Sustained-release formulations of α -glucosidase inhibitors," and incorporated herein by reference in its entirety for all purposes, discloses sustained released formulations of α -glucosidase inhibitors, *e.g.* acarbose useful in the present invention.

Other types of agents that have found limited utility in treating diabetes include potassium channel antagonists such as repaglinide, and aldose reductase inhibitors such as zopolrestat, minalrestat, ponalrestat and tolrestat. Still in the experimental stage, glucagon antagonists, activators of the retinoid-X receptor (RXR), and anti-obesity agents are also being evaluated as potential antidiabetic agents. Others agents suitable for use in the present invention include, but are not limited to, glucagon, and miscellaneous agents such as methyl palmoxirate, palmoxirate sodium, pirogliride, pramlintide, amlintide, seglifide.

10 METHODS, USES, DOSAGES AND SCHEDULES

In another embodiment, the present invention provides a method for modulating conditions associated with diabetes or diabetes-related disorders in a host, comprising administering to the host an efficacious amount of compositions comprising (i) a compound of Formula I in combination with (ii) one or more antidiabetic agents. In certain aspects, the compositions of the present invention that are administered comprise a compound of Formula I formulated together with one or more antidiabetic agents. Alternatively, the composition that is administered comprises a compound of Formula I independently formulated from one or more antidiabetic agents *i.e.*, separately formulated.

The pharmaceutical compositions of the present invention are suitable for use in a variety of drug delivery systems. Examples of suitable formulations for use in the present invention are found in *Remington's Pharmaceutical Sciences* (Mack Publishing Company, Philadelphia, PA, 17th ed. (1985)), which is incorporated herein by reference. In addition, for a brief review of methods for drug delivery, *see*, Langer, *Science* 249:1527-1533 (1990), which is incorporated herein by reference.

The pharmaceutical compositions of the present invention are intended for parenteral, topical, oral or local administration. In certain aspects, the pharmaceutical compositions are administered parenterally, *e.g.*, intravenously, subcutaneously, intradermally, or intramuscularly. In one embodiment, the invention provides compositions for parenteral administration which comprise a compound of Formula I, an antidiabetic agent as described above, dissolved or suspended in an acceptable carrier, preferably an aqueous carrier. A variety of aqueous carriers may be used including, for example, water, buffered water, 0.4% saline, 0.3% glycine, hyaluronic acid and the like. These compositions may be sterilized by conventional, well-known sterilization techniques or, they may be sterile filtered. The resulting aqueous solutions may be packaged for use as is

or lyophilized, the lyophilized preparation being combined with a sterile solution prior to administration. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions including pH adjusting and buffering agents, tonicity adjusting agents, wetting agents and the like, such as, for example, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, sorbitan monolaurate, triethanolamine oleate, *etc.*

For solid formulations, compounds of Formula I can be admixed with conventional nontoxic solid carriers can be used which include, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, talcum, cellulose, glucose, sucrose, magnesium carbonate, and the like. For oral administration, a pharmaceutically acceptable nontoxic composition is formed by incorporating any of the normally employed excipients, such as those carriers previously listed, and generally 10-95% of active ingredient and more preferably at a concentration of 25%-75%.

For aerosol administration, the compounds of Formula I and antidiabetic agents are preferably supplied in finely divided form along with a surfactant and propellant. The surfactant must, of course, be nontoxic, and preferably soluble in the propellant. Representative of such agents are the esters or partial esters of fatty acids containing from 6 to 22 carbon atoms, such as caproic, octanoic, lauric, palmitic, stearic, linoleic, linolenic, olesteric and oleic acids with an aliphatic polyhydric alcohol or its cyclic anhydride. Mixed esters, such as mixed or natural glycerides may be employed. A carrier can also be included, as desired, as with, *e.g.*, lecithin for intranasal delivery.

The compounds of the present invention can be prepared and administered in a wide variety of oral and parenteral dosage forms. Thus, the compounds of the present invention can be administered by injection, that is, intravenously, intramuscularly, intracutaneously, subcutaneously, intraduodenally, or intraperitoneally. Also, the compounds described herein can be administered by inhalation, for example, intranasally. Additionally, the compounds of the present invention can be administered transdermally. Accordingly, the present invention also provides pharmaceutical compositions comprising a pharmaceutically acceptable carrier or excipient and either a compound of Formula I or a pharmaceutically acceptable salt of a compound of Formula I.

For preparing pharmaceutical compositions from the compounds of the present invention, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier can be one or more substances that may also act as

diluents, flavoring agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material.

In powders, the carrier is a finely divided solid that is in a mixture with the finely divided active component. In tablets, the active component is mixed with the carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired.

The powders and tablets preferably contain from 5% to 70% of the active compound. Suitable carriers are magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the formulation of the active compound with encapsulating material as a carrier providing a capsule in which the active component with or without other carriers, is surrounded by a carrier, which is thus in association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges can be used as solid dosage forms suitable for oral administration.

For preparing suppositories, a low melting wax, such as a mixture of fatty acid glycerides or cocoa butter, is first melted and the active component is dispersed homogeneously therein, as by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool, and thereby to solidify.

Liquid form preparations include solutions, suspensions, and emulsions, for example, water or water/propylene glycol solutions. For parenteral injection, liquid preparations can be formulated in solution in aqueous polyethylene glycol solution.

Aqueous solutions suitable for oral use can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizers, and thickening agents as desired. Aqueous suspensions suitable for oral use can be made by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well-known suspending agents.

Also included are solid form preparations that are intended to be converted, shortly before use, to liquid form preparations for oral administration. Such liquid forms include solutions, suspensions, and emulsions. These preparations may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

The pharmaceutical preparation is preferably in unit dosage form. In such form the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

The quantity of active component in a unit dose preparation may be varied or adjusted from 0.1 mg to 1000 mg, preferably 1.0 mg to 100 mg according to the particular application and the potency of the active component. The composition can, if desired, also contain other compatible therapeutic agents.

In therapeutic use for the treatment of obesity, NIDDM, or inflammatory conditions, the compounds utilized in the pharmaceutical method of the invention are administered at the initial dosage of about 0.001 mg/kg to about 100 mg/kg daily. A daily dose range of about 0.1 mg/kg to about 10 mg/kg is preferred. The dosages, however, may be varied depending upon the requirements of the patient, the severity of the condition being treated, and the compound being employed. Determination of the proper dosage for a particular situation is within the skill of the practitioner. Generally, treatment is initiated with smaller dosages that are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under circumstances is reached. For convenience, the total daily dosage may be divided and administered in portions during the day, if desired.

In therapeutic applications, the compounds of Formula I and antidiabetic agents of the present invention are administered to a patient in a combination amount sufficient to elicit a response. An amount adequate to accomplish this is defined as "therapeutically effective combination dose." The methods include the administration of the combination of compound of Formula I with antidiabetic agent wherein the two components are delivered in a simultaneous manner, in combination therapy wherein the compound of Formula I is administered first, followed by the antidiabetic agent, as well as wherein the antidiabetic agent is delivered first followed by the compound of Formula I.

Since the present invention has an aspect that relates to a combination of active ingredients which can be administered separately, the invention also relates to combining separate pharmaceutical compositions in kit form. The kit comprises two separate pharmaceutical compositions: a compound of Formula I, a prodrug thereof or a pharmaceutically acceptable salt and a second compound such as an antidiabetic agent as

described above. The kit comprises a container for containing the separate components such as a divided bottle or a divided foil packet, however, the separate components can also be contained within a single, undivided container. Typically, the kit comprises directions for the administration of the separate components. The kit form is particularly advantageous
5 when the separate components are preferably administered in different dosage forms (*e.g.*, oral and parenteral), are administered at different dosage intervals, or when titration of the individual components of the combination is desired by the prescribing physician.

Effective combination amounts for various uses will depend on, for example, the particular antidiabetic agent, the compound of Formula I employed, the manner of
10 administration, the weight and general state of health of the patient, and the judgment of the prescribing physician. In one embodiment, composition or formulation to be administered will contain a quantity of a compound(s) according to Formula I in an amount effective to treat the disease/condition of the subject being treated, *e.g.*, a glycogen phosphorylase dependent disease/condition. The amount of antidiabetic agent will depend in part to the
15 chemical class.

In certain instances, administration of the compounds of Formula I can be via any method which provides systemic exposure to the compound of this invention, preferably to the muscle and fatty tissue. These methods include oral routes, parenteral, intraduodenal routes, *etc.* Generally, the compounds of the present invention are
20 administered in single (*e.g.*, once daily) or multiple doses. The compounds of the present invention are generally administered in the form of a pharmaceutical composition comprising at least one of the compounds of Formula I together with a pharmaceutically acceptable carrier or diluent. Thus, the compounds of this invention can be administered individually or together in any conventional oral, parenteral or transdermal dosage form.

For oral administration, a pharmaceutical composition can take the form of
25 solutions, suspensions, tablets, pills, capsules, powders, and the like. Tablets containing various excipients such as sodium citrate, calcium carbonate and calcium phosphate are employed along with various binders such as starch and preferably potato or tapioca starch and certain complex silicates, together with binding agents such as polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate,
30 sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type are also employed as fillers in soft and hard-filled gelatin capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are

desired for oral administration, the compounds of this invention can be combined with various sweetening agents, flavoring agents, coloring agents, emulsifying agents and/or suspending agents, as well as such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

5 It will be appreciated that the actual preferred course of therapy will vary according to, *inter alia*, the mode of administration of the compound of Formula I, the particular formulation of the antidiabetic agent being utilized, the mode of administration of the compounds, the particular disease being treated and the particular host being treated. The optimal course of therapy for a given set of conditions can be ascertained by those
10 skilled in the art using conventional course of therapy determination tests and in view of the information set out herein.

ADDITIONAL USES FOR COMPOUNDS OF FORMULA I

 The compounds of Formula I have been recently shown to interact with
15 PPAR γ from human and other species. Some of the compounds of Formula I show various degrees of intrinsic activity towards PPAR γ ; that is, some of them display significant PPAR γ activation, while others possess a distinct PPAR γ antagonist-like profile. As such, specific compounds of Formula I are well suited for the treatment of the various conditions and diseases recently established to be mediated by, or linked to
20 PPAR γ activity levels. Tables I, II, and III below outline other indications in which the compositions of the present invention are useful.

TABLE IMetabolic Conditions**1. Diabetes and Conditions Secondary to Diabetes**

- Hypertension
- Angina pectoris
- Dyslipidemia
 - Hypertriglyceridemia
 - Hyperlipoproteinemias
 - Hypercholesterolemia
- Gout
- Nephropathy and other renal diseases secondary to diabetes
- Diabetic neuropathy.

2. Other insulin-resistance-related diseases

- Polycystic ovarian syndrome,
- Glucocorticoid-induced insulin resistance

3. Obesity

• PPAR γ agonists promote adipocyte differentiation and fat accumulation. PPAR γ antagonists block normal hormone-mediated differentiation of preadipocytes into adipocytes. (See, Wright *et al.*, *J. Biol. Chem.* **275**(3) 1873-1877 (2000).)

• PPAR γ agonists inhibit expression of ob gene (leptin production) in mature adipocytes. It follows that PPAR γ antagonists will increase leptin production with ensuing decrease in appetite and food consumption. (See, Sinha *et al.*, *Metab. Clin. Exp.*, **48** (6) 786-791 (1999).)

• Hyperleptinemia condition (which can be induced with a PPAR γ antagonist) in rats results in downregulation of PPAR γ expression and upregulation of fatty acid-oxidizing enzymes. Also accompanied by a reversal of adipocyte differentiation. Suggests that relatively short-term treatment with high dose of PPAR γ antagonist will have long-lasting effects on obesity (conventional treatments of obesity reduce fat content in mature

adipocytes but leaves them with lipogenic enzymes capable of rapid resynthesis of fat, likely to be responsible for treatment failure). (See, Zhou *et al.*, *Proc. Natl. Acad. Sci. USA* **96** (5) 2391-2395 (1999).)

- PPAR γ agonists upregulate UCP2 expression in adipocytes and skeletal muscle, resulting in increased energy expenditure. (See, Viguerie-Bascands *et al.*, *Biochem. Biophys. Res. Commun.* **256** (1) 138-141 (1999) and Camirand *et al.*, *Endocrinology* **139** (1) 428-431 (1998).)

- PPAR γ is critical for controlling expression of UCP2 and UCP3 in adipose tissue. (See, Kelly *et al.*, *Endocrinology* **139** (12) 4920-4927 (1998).)

4. Hypertension

- PPAR γ agonists suppress endothelin-1 secretion by vascular endothelial cells and result in decreased blood pressure. (See, Satoh *et al.*, *Biochem. Biophys. Res. Commun.* **254** (3) 757-763 (1999) and Itoh *et al.*, *Clin. Exp. Pharmacol. Physiol.* **26** (7) 558-560 (1999).)

- PPAR γ agonists decrease blood pressure in various models of hypertension. (See, Komers *et al.*, *Physiol. Res. (Prague)* **47** (4) 215-225 (1998).)

5. Lipid Disorders

- PPAR γ has been implicated in systemic glucose and lipid homeostasis. (See, Kliewer *et al.*, *Curr. Opin. Genet. Dev.* **8** (5) 576-581 (1998).)

- PPAR γ agonists improve hypertriglyceridemia. (See, Berger *et al.*, *J. Biol. Chem.* **274** (10) 6718-6725 (1999).)

- PPAR γ agonists are antihyperlipidemic. (See, Henke *et al.*, *J. Med. Chem.* **41** (25) 5020-5036 (1998).)

- PPAR γ activator shown to increase high-density lipoprotein (HDL) in dose-dependent manner. Also decreased VLDL, LDL and triglycerides. (See, Bisgaier *et al.*, *J. Lipid Res.* **39** (1) 17-30 (1998).)

6. Atherosclerosis

- Activated monocytes/macrophages express PPAR γ and PPAR γ activation downregulates the induced macrophage production of IL-1 and TNF α . Potential implications for atherosclerosis. (See, McCarty *et al.*, *J. Med. Food* 1 (3) 217-226 (1999).)
- PPAR γ mediates the effects of non-esterified fatty acids (NEFA) on smooth muscle cells, which alter the extracellular matrix in the intima of small and large arteries. These changes can lead to increased deposition of LDL and may be associated with the etiology of atherosclerosis. Modulators of PPAR γ will affect this process. (See, Olsson *et al.*, *Diabetes* 48 (3) 616-622 (1999).)
- PPAR γ agonists inhibit proliferation, hypertrophy and migration of vascular smooth muscle cells induced by growth factors. These processes are crucial in the development of vascular remodeling, atherosclerosis.
- PPAR γ involved in negative regulation of monocyte/macrophage function in atherosclerotic plaques and regulates expression of matrix metalloprotease-9, an enzyme implicated in plaque rupture. In this case, a PPAR γ agonist may be useful. (See, Marx *et al.*, *Am. J. Pathol.* 153 (1) 17-23 (1998) and Shu *et al.*, *Biochem. Biophys. Res. Commun.* 267 (1) 345-349 (2000).)
- PPAR γ is expressed in macrophage foam cells of human sclerotic lesions. (See, Ricote *et al.*, *Proc. Natl. Acad. Sci. USA* 95 (13) 7614-7619 (1998).)
- PPAR γ is expressed in atherosclerotic plaques and in endothelial cells. In endothelial cells, PPAR γ agonists markedly attenuate the TNF α -induced expression of VCAM-1 and ICAM-1 (vascular cell adhesion molecules) in vitro. PPAR γ agonists significantly reduce monocyte/macrophage homing to atherosclerotic plaques in apoE-deficient mice. These effects combined may have beneficial effects in modulating inflammatory response in atherosclerosis. (See, Pasceri *et al.*, *Circulation* 101 (3) 235-238 (2000).)

- Human genetic evidence also suggests that PPAR γ plays significant role in atherogenesis, independently from effects on obesity and lipid metabolism, possibly via a direct local vascular wall effect. (See, Wang *et al.*, *Cardiovasc. Res.* **44** (3) 588-594 (1999).)

- In the past year, there has been a very significant increase in research implicating PPAR γ in macrophage biology, cell cycle regulation and atherosclerosis. PPAR γ as regulator of monocyte/macrophage function. (See, Ricote *et al.*, *J. Leukocyte Biol.* **66** (5) 733-739 (1999).)

7. Bone Disorders

- TZDs inhibit in vitro bone nodule formation and mineralization. (See, Johnson *et al.*, *Endocrinology* **140** (7) 3245-3254 (1999).)

- PPAR γ polymorphism affects bone mineral density in postmenopausal women. (See, Ogawa *et al.*, *Biochem. Biophys. Res. Commun.* **260** (1) 122-126 (1999).)

- TZDs are potent inhibitors of bone resorption in vitro. Thus, TZDs may suppress bone resorption in diabetic patients and prevent bone loss. (See, Okazaki *et al.*, *Endocrinology* **140** (11) 5060-5065 (1999).)

- Short-term treatment of diabetic patients with TZD decreases bone turnover. This effect is noted before significantly improvement on glucose metabolism, suggesting that effect is direct on bone. Dual effects on glucose and bone metabolism may lead to spared bone mass in diabetic patients. (See, Okazaki *et al.*, *Endocr. J. (Tokyo)* **46** (6) 795-801 (1999).)

8. Female – Specific Conditions

- PPAR γ agonists can be used to inhibit excessive uterine bleeding in women (climacteric). (See, Urban *et al.*, WO-9839006.)

9. Acne

- PPAR γ implicated in the differentiation of sebocytes. PPAR γ agonists may be used in the treatment of acne. (See, Rosenfield *et al.*, *Dermatology* (Basel) **196** (1) 43-46 (1998).)
- Other skin disorders associated with differentiation of epidermal cells. (See, Rivier *et al.*, FR 2773075-A1.)
- Proliferative diseases of the skin. (See, Pershadsingh *et al.*, US 5,981,586.)

10. Cell Proliferation

- In combination with a retinoid-X receptor agonist, a PPAR γ agonist reduces uncontrolled cell proliferation, including cancer, restenosis and atherosclerosis. PPAR γ agonists, alone or in combination with known agents, may reduce proliferative response seen following angioplasty, vessel transplant or endarectomy.

TABLE 2**CNS****1. Alzheimer's**

- PPAR γ agonists inhibit b-amyloid stimulated secretion of proinflammatory products by microglia and monocytes that are responsible for neurotoxicity and astrocyte activation. They also arrest differentiation of monocytes into activated macrophages, and inhibit b-amyloid-stimulated expression of IL-6, TNF α and cyclooxygenase-2. (See, Combs *et al.*, *J. Neuroscience* **20** (2) 558-567 (2000).)
- In temporal cortex from AD patients, cyclooxygenase-1 and -2 levels were increased. PPAR γ levels were also increased. Certain agents that activate PPAR γ inhibit COX-2 expression in glial cells. (See, Kitamura *et al.*, *Biochem. Biophys. Res. Commun.* **254** (3) 582-586 (1999).)
- PPAR γ agonists protect cerebellar granule cells from cytokine-induced apoptotic death

by inhibition of iNOS. (See, Heneka *et al.*, *J. Neuroimmunol.* **100** (1-2) 156-168 (1999).)

- Activated monocytes/macrophages express PPAR γ and PPAR γ activation downregulates the induced macrophage production of IL-1 and TNF α . Potential implications for AD. (See, McCarty *et al.*, *J. Med. Food* **1** (3) 217-226 (1999).)

2. Neuroinflammation

- PPAR γ agonists inhibit LPS and IFN-g induced expression of iNOS by glial cells. (See, Kitamura *et al.*, *Neurosci. Lett.* **262** (2) 129-132 (1999).)

- PPAR γ ligands may also be relevant for other disorders associated with neuroinflammation, such as ischemic stroke, closed-head injury, and multiple sclerosis.

TABLE 3

Chemotherapy

1. Cancer

- Antiangiogenic affect of PPAR γ agonists are mediated by apoptotic stimulus on endothelial cells. (See, Bishop-Balley *et al.*, *J. Biol. Chem.* **274** (24) 17042-17048 (1999).)

- PPAR γ agonists induce terminal differentiation and growth arrest of human colon cancer cells. (See, Kitamura *et al.*, *Jpn. J. Cancer Res.* **90** (1) 75-80 (1999) and Sarraf *et al.*, *Nat. Med.* (NY) **4** (9) 1046-1052 (1998).)

- PPAR γ agonists enhance the antiproliferative effects of retinoic acid on human colon cancer cells. (See, Brockman *et al.*, *Gastroenterology* **115** (5) 1049-1055 (1998).)

- PPAR γ agonist has potent antitumor effect against human prostate cancer in vitro and in vivo. (See, Kubota *et al.*, *Cancer Res.* **58** (15) 3344-3352 (1998).)

- PPAR γ agonists cause inhibition of proliferation of cultured human breast tumor cells and induces apoptosis. Effects also seen in vivo (mice). (See, Elstner *et al.*, *Proc. Natl.*

Acad. Sci. USA **95** (15) 8806-8811 (1998).)

- PPAR γ agonists induce terminal differentiation of malignant breast epithelial cells. (See, Mueller *et al.*, *Mol. Cell* **1** (3) 465-470 (1998) and Yee *et al.*, *Int. J. Oncol.* **15** (5) 967-973 (1999).)
- PPAR γ agonists are useful in the treatment of liposarcomas. (See, Evans *et al.*, WO-9829120.)
- PPAR γ highly expressed in all human transitional epithelial cell cancers, including uroepithelial human cancers. PPAR γ agonists induce differentiation and inhibit proliferation. (See, Guan *et al.*, *Neoplasia* (NY) **1** (4) 330-339 (1999).)
- Differentiation of many cell types (hepatocytes, fibroblasts, adipocytes, keratinocytes, myocytes, and monocyte/macrophages) involves PPAR γ . PPAR γ modulators may play role in treating malignancies that result from these and other cell types. (See, Varmecq *et al.*, *Lancet* **354** (9173) 141-148 (1999).)

TABLE 4

Inflammation / Immune

1. Inflammation / Immune

- PPAR γ is markedly upregulated in activated macrophages. PPAR γ involved in negative regulation of monocyte/macrophage function, including generation of inflammatory cytokines, expression of iNOS, gelatinase B and scavenger receptor A. PPAR γ agonists may be of value. (See, Marx *et al.*, *Am. J. Pathol.* **153** (1) 17-23 (1998).)
- Incremental therapeutic benefit of NSAIDs (some of which activate PPAR γ) in the treatment of rheumatoid arthritis may be mediated via PPAR γ activation. (See, Jiang *et al.*, *Nature* **391** (6662) 82-86 (1998).)
- PPAR γ agonists inhibit iNOS production by activated macrophages. PPAR γ agonists may be useful. (See, Colville-Nash *et al.*, *J. Immunol.* **161** (2) 978-984 (1998).)

- PPAR γ agonists attenuate antigen-induced cytokine production by bone-marrow-derived mast cells. (See, Sugiyama *et al.*, *FEBS Lett.* **467** (2-3) 259-262 (2000).)
- Recently, an immunomodulatory role for PPAR γ has been described in cells critical to the innate immune system, the monocyte/macrophage. PPAR γ agonists have been shown to mediate significant inhibition of proliferative responses of helper T-cell clones and freshly isolated splenocytes. Thus, IL-2 production by the T-cell clones is inhibited by PPAR γ agonists, which may have utility as immunosuppressants. (See, Clark *et al.*, *J. Immunol.* **164** (3) 1364-1371 (2000).)
- PPAR γ as regulator of monocyte/macrophage function. (See, Ricote *et al.*, *J. Leukocyte Biol.* **66** (5) 733-739 (1999).)
- PPAR γ expression in WBC may play a role in host response to acute inflammatory challenge and may prove to be an important target for anti-inflammatory control. (See, Leininger *et al.*, *Biochem. Biophys. Res. Commun.* **263** (3) 749-753 (1999).)
- PPAR γ activators may help limit chronic inflammation mediated by vascular cell adhesion molecule VCAM-1 and monocytes. (See, Jackson *et al.*, *Arterioscler., Thromb., Vasc. Biol.* **19** (9) 2094-2104 (1999).)
- PPAR γ agonists significantly attenuate cytokine production in colon cell lines by inhibiting activation of the nuclear factor NF-kB. PPAR γ agonists also markedly reduce colonic inflammation in a mouse model of inflammatory bowel disease (IBD). PPAR γ agonists may be useful in treating colitis and Crohn's disease (human forms of IBD). (See, Su *et al.*, *J. Clin. Invest.* **104** (4) 383-389 (1999).)

2. Ophthalmic**Macular Degeneration**

- Antiangiogenic affect of PPAR γ agonists mediated by apoptotic stimulus on endothelial cells. (See, Bishop-Balley *et al.*, *J. Biol. Chem.* (1999) **274** (24) 17042-17048.)

3. Antiangiogenic

The invention will be described in greater detail by way of specific examples.


The following examples are offered for illustrative purposes, and are not intended to limit the invention in any manner.

5

EXAMPLES**Example 1**

This example sets forth a synthesis procedure used to generate compounds of Formula I. Compounds of Formula I were synthesized using methods as outlined in

10 Figure 1.




To a mixture of 5-amino-2-(3-chloro-5-pyridyloxy)benzonitrile (0.457 g) in methylene chloride was added 2,4-dichlorobenzenesulfonyl chloride (0.456 g, from
15 Maybridge), followed by pyridine (150 μ L). The reaction progress was monitored by TLC, and upon completion the solvent was removed under vacuum. The resulting residue was partitioned between methylene chloride and water. The organic layer was drawn off and concentrated. The residue was triturated with ether to provide 0.447 g of the title compound as a white solid, mp 154-156 $^{\circ}$ C.

20 ^1H NMR (400 MHz) (CDCl_3) δ 8.59 (s, 1H); 8.42 (s, 1H) 8.08 (d, $J=8.5$ Hz, 1H); 7.72(t, $J=1.8$, 1H); 7.605 (d, $J=2.7$ Hz, 1H) 7.53 (dd, $J=8.5$, 2 Hz, 1H); 7.48 (dd, $J=9.4$ Hz, 1H); 7.22 (s, 1H); 7.0 (d, $J=9.0$ Hz, 1H). m/e (M-H) 456.

The product from above was oxidized to the corresponding pyridine N-oxide using 3-chloroperoxybenzoic acid in methylene chloride to provide the product as a white solid. m/e 470 ($M+H$).

Using a similar method and protocol, the following compounds were
5 generated.



Various other compounds of Formula I are illustrated in Figure 3.

10

Example 2

This example illustrates combination therapy of a compound of Formula I and a glitazone by oral administration.

Patients having NIDDM (Type II diabetes mellitus) are selected for
15 therapy. The patients weigh between 70-100 kilograms. A compound of Formula I is orally administered in a dosage of 20 to 1,000 milligrams twice daily, more typically 100 mg twice daily. For infants or children the doses suggested are lowered in a linear fashion based on body weight or surface area.


Half the patient population is administered troglitazone as well as a compound of Formula I using an effective dose of both agents. The other half of the patients are administered an effective dose of troglitazone. The patients are monitored for improvement in the manifestations of the disease and for side effects, such as body weight gain and signs of liver toxicity.

The results indicate the administration of a combination of i) a compound of Formula I with ii) troglitazone increases the efficacy of either agent alone. The composition also provides concomitant decrease in the side effects of either agent alone.

All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

WHAT IS CLAIMED IS:

- 1 1. A pharmaceutical composition, said composition comprising:
 2 (i) a compound having the formula I or a pharmaceutically acceptable salt
 3 thereof:



4 wherein

5 Ar¹ is an aryl group;

6 X is a divalent linkage selected from the group consisting of (C₁-
 7 C₆)alkylene, (C₁-C₆)alkylenoxy, (C₁-C₆)alkylenamino, (C₁-C₆)alkylene-S(O)_k-, -O-, -
 8 C(O)-, -N(R¹¹)-, -N(R¹¹)C(O)-, -S(O)_k- and a single bond,

9 wherein

10 R¹¹ is a member selected from the group consisting of hydrogen, (C₁-
 11 C₈)alkyl, (C₁-C₈)heteroalkyl and aryl(C₁-C₄)alkyl; and the subscript k is an integer of
 12 from 0 to 2;

13 Y is a divalent linkage selected from the group consisting of alkylene, -O-,
 14 -C(O)-, -N(R¹²)-S(O)_m-, -N(R¹²)-S(O)_m-N(R¹³)-, -N(R¹²)C(O)-, -S(O)_n- and a single bond,
 15 wherein

16 R¹² and R¹³ are members independently selected from the group consisting
 17 of hydrogen, (C₁-C₈)alkyl, (C₁-C₈)heteroalkyl and aryl(C₁-C₄)alkyl; and the subscripts m
 18 and n are independently integers of from 0 to 2;

19 R¹ is a member selected from the group consisting of hydrogen,
 20 heteroalkyl, aryl, arylalkyl, halogen, cyano, nitro, (C₁-C₈)alkyl, (C₁-C₈)alkoxy, -C(O)R¹⁴,
 21 -CO₂R¹⁴, -C(O)NR¹⁵R¹⁶, -S(O)_p-R¹⁴, -S(O)_q-NR¹⁵R¹⁶, -O-C(O)-OR¹⁷, -O-C(O)-R¹⁷, -O-
 22 C(O)-NR¹⁵R¹⁶, -N(R¹⁴)-C(O)-NR¹⁵R¹⁶, -N(R¹⁴)-C(O)-R¹⁷ and -N(R¹⁴)-C(O)-OR¹⁷;

23 wherein

24 R¹⁴ is a member selected from the group consisting of hydrogen, (C₁-
 25 C₈)alkyl, (C₁-C₈)heteroalkyl, aryl and aryl(C₁-C₄)alkyl;

R^{15} and R^{16} are members independently selected from the group consisting of hydrogen, (C₁-C₈)alkyl, (C₁-C₈)heteroalkyl, aryl, and aryl(C₁-C₄)alkyl, or taken together with the nitrogen to which each is attached form a 5-, 6- or 7-membered ring;

R^{17} is a member selected from the group consisting of alkyl, heteroalkyl, aryl and arylalkyl;

the subscript p is an integer of from 0 to 3; and

the subscript q is an integer of from 1 to 2; and

R^2 is a member selected from the group consisting of (C₁-C₈)alkyl, (C₁-C₈)heteroalkyl, aryl and aryl(C₁-C₄)alkyl; and

R^3 is a member selected from the group consisting of halogen, cyano, nitro, (C₁-C₈)alkyl and (C₁-C₈)alkoxy; and

ii) one or more antidiabetic agents, prodrugs thereof, or pharmaceutically acceptable salts of said antidiabetic agent; and optionally a pharmaceutically acceptable carrier or diluent.

2. The pharmaceutical composition of claim 1, wherein said antidiabetic agent is a member selected from the group consisting of a sulfonylurea, a biguanide, a glitazone, a PPAR γ modulator, an α -glucosidase inhibitor, a potassium channel antagonist, an aldose reductase inhibitor, a glucagon antagonist, an activator of RXR, insulin therapy, an anti-obesity agent and mixtures thereof.

3. The pharmaceutical composition of claim 2, wherein said sulfonyl urea is a member selected from the group consisting of acetohexamide, chlorpropamide, glyburide, glipizide, gliclazide, glimepiride, gliquidone, glisoxepid, glibornuride, gliamilide, glicetanile, gliflumide, glymidine, glyparamide, tolpyrramide, glyhexamide, phenbutamide, tolazamide, tolbutamide and tolcyclamide.

4. The pharmaceutical composition of claim 2, wherein said biguanide is a member selected from the group consisting of metformin, buformin, etoformin and phenformin.

5. The pharmaceutical composition of claim 2, wherein said glitazone is a member selected from the group consisting of troglitazone, rosiglitazone, pioglitazone, JTT-501, YM-440, ciglitazone, darglitazone, englitazone, AD-5075 and BM-131246.

1 6. The pharmaceutical composition of claim 2, wherein said α -
2 glucosidase inhibitor is a member selected from the group consisting of acarbose,
3 miglitol, emiglitate, voglibose, MDL-25,637, camiglibose and MDL-73,945.

1 7. The pharmaceutical composition of claim 1, wherein
2 X is selected from the group consisting of -O-, -S(O)_k-, -N(R¹¹)-, -CH₂-,
3 -C(O)- and a single bond;
4 Y is -N(R¹²)-S(O)_m-;
5 wherein
6 R¹¹ is hydrogen or (C₁-C₈) alkyl;
7 R¹² is selected from the group consisting of hydrogen and (C₁-C₈)alkyl;
8 and the subscript m is an integer of from 0 to 2;
9 R¹ is a member selected from the group consisting of hydrogen, halogen,
10 cyano, (C₁-C₈)alkoxy, (C₁-C₈)alkyl, -C(O)R¹⁴, -CO₂R¹⁴ and -C(O)NR¹⁵R¹⁶;
11 wherein
12 R¹⁴ is a member selected from the group consisting of hydrogen, (C₁-
13 C₈)alkyl, (C₁-C₈)heteroalkyl, aryl and aryl(C₁-C₄)alkyl;
14 R¹⁵ and R¹⁶ are members independently selected from the group consisting
15 of hydrogen, (C₁-C₈)alkyl, (C₁-C₈)heteroalkyl, aryl, and aryl(C₁-C₄)alkyl, or taken
16 together with the nitrogen to which each is attached form a 5-, 6- or 7-membered ring;
17 R² is a member selected from the group consisting of aryl and aryl(C₁-
18 C₄)alkyl; and
19 R³ is a member selected from the group consisting of halogen, cyano, (C₁-
20 C₈)alkyl and (C₁-C₈)alkoxy.

1 8. The pharmaceutical composition of claim 7, wherein Ar¹ is a
2 substituted or unsubstituted heteroaryl group having from 1 to 3 heteroatoms as ring
3 members.

1 9. The pharmaceutical composition of claim 7, wherein Ar¹ is a
2 phenyl, naphthyl, benzothiazole, benzimidazole, quinoline, isoquinoline or group having
3 from 1 to 3 substituents selected from the group consisting of halogen, cyano, CF₃, nitro,
4 (C₁-C₈)alkyl and (C₁-C₈)alkoxy.

1 10. The pharmaceutical composition of claim 1, wherein X is a
2 divalent linkage selected from the group consisting of -O-, -C(O)-, N(R¹¹)-, -CH₂-,
3 -S(O)_k- and a single bond.


1 11. The pharmaceutical composition of claim 1, where Y is a divalent
2 linkage -N(R¹²)-S(O)₂-; and R¹² is hydrogen or (C₁-C₈)alkyl, and R² is a member selected
3 from the group consisting of aryl and arylalkyl.

1 12. The pharmaceutical composition of claim 1, wherein R¹ is selected
2 from the group consisting of hydrogen, halogen, cyano, (C₁-C₈)alkoxy, (C₁-C₈)alkyl, -
3 C(O)R¹⁴, -CO₂R¹⁴ and -C(O)NR¹⁵R¹⁶ wherein R¹⁴ is (C₁-C₈)alkyl, and R¹⁵ and R¹⁶ are
4 independently selected from the group consisting of hydrogen and (C₁-C₈)alkyl, or taken
5 together with the nitrogen to which each is attached form a 5- or 6-membered ring.


1 13. The pharmaceutical composition of claim 1, wherein Ar¹ is
2 substituted with from one to two substituents selected from the group consisting of
3 halogen, -OCF₃, -OH, -O(C₁-C₆)alkyl, -CF₃, (C₁-C₈)alkyl and -NO₂.

1 14. The pharmaceutical composition of claim 1, wherein Ar¹ is a
2 mono- or bicycle heteroaryl group having from 1 to 2 nitrogen atoms as ring members.


1 15. The pharmaceutical composition of claim 1, wherein said
2 compound is represented by a formula selected from the group consisting of




(Ia)




(Ib)




(Ic)




(Id)




(Ie)



(If)




(Ig)



(Ih)

and



(Ii)


16. The pharmaceutical composition of claim 1, wherein X is a divalent linkage selected from the group consisting of -O-, -S-, -S(O)-, -S(O)₂-, -NH-, -CH₂-, -C(O)- and a single bond.

17. The pharmaceutical composition of claim 1, wherein Y is -NH-S(O)₂- or -NH-S(O)₂-NH-, and R² is aryl.

18. The pharmaceutical composition of claim 1, wherein R² is selected from the group consisting of substituted phenyl, substituted pyridyl and substituted naphthyl, wherein the substituents number from one to three and are independently selected from the group consisting of halogen, -OCF₃, -OH, -O(C₁-C₈)alkyl, -C(O)-(C₁-C₈)alkyl, -CN, -CF₃, (C₁-C₈)alkyl and -NH₂.


19. The pharmaceutical composition of claim 1, wherein R¹ is selected from the group consisting of -H, halogen, C(O)R¹⁴, -(C₁-C₈)alkyl, and -CONR¹⁵R¹⁶ wherein R¹⁵ and R¹⁶ are each independently selected from hydrogen, alkyl, aryl and arylalkyl, and R¹⁴ is (C₁-C₈) alkyl.

20. The pharmaceutical composition of claim 1, wherein said compound of Formula I is selected from the group consisting of



21. A method for modulating medical conditions associated with metabolic disorders, said method comprising administering to said host an efficacious amount of

(i) a compound having the formula:



wherein

Ar^1 is an aryl group;

X is a divalent linkage selected from the group consisting of (C₁-C₆)alkylene, (C₁-C₆)alkylenoxy, (C₁-C₆)alkylenamino, (C₁-C₆)alkylene-S(O)_k-, -O-, -C(O)-, -N(R¹¹)-, -N(R¹¹)C(O)-, -S(O)_k- and a single bond,

wherein

R¹¹ is a member selected from the group consisting of hydrogen, (C₁-C₈)alkyl, (C₁-C₈)heteroalkyl and aryl(C₁-C₄)alkyl; and the subscript k is an integer of from 0 to 2;

Y is a divalent linkage selected from the group consisting of alkylene, -O-, -C(O)-, -N(R¹²)-S(O)_m-, -N(R¹²)-S(O)_m-N(R¹³)-, -N(R¹²)C(O)-, -S(O)_n- and a single bond,

wherein

R¹² and R¹³ are members independently selected from the group consisting of hydrogen, (C₁-C₈)alkyl, (C₁-C₈)heteroalkyl and aryl(C₁-C₄)alkyl; and the subscripts m and n are independently integers of from 0 to 2;

R¹ is a member selected from the group consisting of hydrogen, heteroalkyl, aryl, arylalkyl, halogen, cyano, nitro, (C₁-C₈)alkyl, (C₁-C₈)alkoxy, -C(O)R¹⁴, -CO₂R¹⁴, -C(O)NR¹⁵R¹⁶, -S(O)_p-R¹⁴, -S(O)_q-NR¹⁵R¹⁶, -O-C(O)-OR¹⁷, -O-C(O)-R¹⁷, -O-C(O)-NR¹⁵R¹⁶, -N(R¹⁴)-C(O)-NR¹⁵R¹⁶, -N(R¹⁴)-C(O)-R¹⁷ and -N(R¹⁴)-C(O)-OR¹⁷;

wherein

R¹⁴ is a member selected from the group consisting of hydrogen, (C₁-C₈)alkyl, (C₁-C₈)heteroalkyl, aryl and aryl(C₁-C₄)alkyl;

R¹⁵ and R¹⁶ are members independently selected from the group consisting of hydrogen, (C₁-C₈)alkyl, (C₁-C₈)heteroalkyl, aryl, and aryl(C₁-C₄)alkyl, or taken together with the nitrogen to which each is attached form a 5-, 6- or 7-membered ring;

R¹⁷ is a member selected from the group consisting of alkyl, heteroalkyl, aryl and arylalkyl;

the subscript p is an integer of from 0 to 3; and

the subscript q is an integer of from 1 to 2; and

R² is a member selected from the group consisting of (C₁-C₈)alkyl, (C₁-C₈)heteroalkyl, aryl and aryl(C₁-C₄)alkyl; and

R³ is a member selected from the group consisting of halogen, cyano, nitro, (C₁-C₈)alkyl and (C₁-C₈)alkoxy; and

ii) at least one therapeutically active agents or a prodrug thereof, or a pharmaceutically acceptable salt of said agent, and a pharmaceutically acceptable carrier or diluent.

22. The method of claim 21, wherein said medical condition is a member selected from the group consisting of diabetes, hypertension, angina, dyslipidemias, gout, diabetic neuropathy, obesity, polycystic ovarian syndrome, hypertriglyceridemia, hypercholesterolemia, atherosclerosis, bone disorders and acne, proliferative diseases, cancer, vascular restenosis, CNS disorders, Alzheimer's disease neuroinflammation, inflammatory disorders, ophthalmic disorders, and reproductive system disorders,

23. The method of claim 21, wherein said compound of Formula I and said at least one therapeutically active agent are formulated together.

24. The method of claim 21, where said compound of Formula I and said at least one therapeutically active agent are formulated separately.

25. The method of claim 21, wherein said medical condition is diabetes, and said therapeutically active agent is an antidiabetic agent.

26. The method of claim 21, wherein said therapeutically active agent is selected from the group consisting of a sulfonylurea, a biguanide, a glitazone, a PPAR γ modulator, an α -glucosidase inhibitor, a potassium channel antagonist, an aldose

4 reductase inhibitor, a glucagon antagonist, an activator of RXR, insulin therapy, an anti-
5 obesity agent, and mixtures thereof.

1 **27.** The method of claim 21, wherein said medical condition is
2 hypertension, and said therapeutically active agent is selected from the group consisting
3 of α_2 -adrenergic agonists, α_1 -adrenergic antagonists, β -adrenergic antagonists, calcium
4 blockers, ACE inhibitors, ATII receptor blockers and vasodilators.

1 **28.** The method of claim 27, wherein said therapeutically active agent
2 is selected from the group consisting of clonidine, prazosin, propranolol, verapamil,
3 captopril, losartan and hydralazine.

1 **29.** The method of claim 21, wherein said medical condition is gout,
2 and said therapeutically active agent is selected from the group consisting of colchicine
3 and allopurinol.

1 **30.** The method of claim 21, wherein said medical condition is obesity,
2 and said therapeutically active agent is a stimulant selected from the group consisting of
3 amphetamine and sibutramine.

1 **31.** The method of claim 21, wherein said medical condition is a
2 member selected from the group consisting of hypertriglyceridemia and
3 hypercholesterolemia and said therapeutically active agent is selected from the group
4 consisting of bile-acid sequestrants, HMG-CoA inhibitors, niacin, and fibrates.

1 **32.** The method of claim 31, wherein said therapeutically active agent
2 is selected from the group consisting of colestipol, simvastatin, niacin, and gemfibrozil.

1 **33.** The method of claim 21, wherein said medical condition is angina,
2 and said therapeutically active agent is selected from the group consisting of
3 nitroglycerine, β -adrenergic blockers and calcium blockers.

1 **34.** The method of claim 33, wherein said therapeutically active agent
2 is selected from the group consisting of nitroglycerine, propranolol, diltiazem and
3 verapamil.

1 **35.** The method of claim 21, wherein said medical condition is
2 atherosclerosis, and said therapeutically active agents is selected from the group
3 consisting of HMG-CoA inhibitors, bile-acid sequestrants and fibrates.

1 **36.** The method of claim 35, wherein said therapeutically active agent
2 is selected from the group consisting of simvastatin, colesteralamine and gemfibrozil.

1 **37.** The method of claim 21, wherein said medical condition are bone
2 disorders/osteoporosis, and said therapeutically active agent is selected from the group
3 consisting of estrogen supplements, and bisphosphonates.

1 **38.** The method of claim 37, wherein said therapeutically active agent
2 is alendronate.

1 **39.** The method of claim 21, wherein said medical condition is acne,
2 and said therapeutically active agent is selected from the group consisting of antibiotics,
3 isotretinoin, and steroid anti-inflammatories.

1 **40.** The method of claim 39, wherein said therapeutically active agent
2 is selected from the group consisting of erythromycin and triamcinolone.

1 **41.** The method of claim 21, wherein said medical condition is cancer,
2 and said therapeutically active agent is selected from the group consisting of alkylating
3 drugs, antimetabolites, microtubule inhibitors, podophyllotoxins, antibiotics, nitrosoureas
4 metal derivatives, hormone therapies and biological therapies.

1 **42.** The method of claim 41, wherein said therapeutically active agent
2 is selected from the group consisting of cyclophosphamide, methotrexate, paclitaxel,
3 etoposide, doxorubicin, carmustine, cisplatin, tamoxifen and Interferon.

1 **43.** The method of claim 21, wherein said medical condition is
2 Alzheimer's disease, and said therapeutically active agent is selected from the group
3 consisting of acetylcholinesterase inhibitors, estrogen therapy and NSAIDs.

1 **44.** The method of claim 43, wherein said therapeutically active agent
2 is selected from the group consisting of donepezil and indomethacin.

1 **45.** The method of claim 21, wherein said medical condition is
2 neuroinflammation, and said therapeutically active agent is selected from the group
3 consisting of NSAIDs, aspirin, vitamin E, and inhibitors of neuronal nitric oxide synthase.

1 **46.** The method of claim 45, wherein said therapeutically active agents
2 is indomethacin.

1 **47.** The method of claim 21, wherein said medical condition is an
2 immune-system dysfunction, and said therapeutically active agent is selected from the
3 group consisting of corticosteroids, interferon- β , cyclosporine, FK-506 and methotrexate.

1 **48.** The method of claim 47, wherein said therapeutically active agent
2 is prednisone.

1 **49.** The method of claim 21, wherein said medical condition is
2 ophthalmic disorders associated with neovascularization and said therapeutically active
3 agent is selected from the group consisting of visodyne, laser phototherapy and VEGF
4 antibodies.

1 **50.** The method of claim 21, wherein said medical condition is
2 vascular restenosis, and such therapeutically active agent is selected from the group
3 consisting of β -adrenergic antagonists, Calcium blockers, aspirin and ACE inhibitors.

1 **51.** The method of claim 50, wherein said therapeutically active agent
2 is selected from the group consisting of propranolol, verapamil and captopril.

1/7

FIGURE 1















FIGURE 2

PPAR γ Agonists


1.		TROGLITAZONE
2.		ROSIGLITAZONE (BRL4963)
3.		PIOGLITAZONE
4.		JTT 501
5.		YM 440 (YAMANOUCHI)
6.		CIGLITAZONE
7.		DARGLITAZONE
8.		ENGLITAZONE
9.		AD-5075 (MERCK)
10.		BM 131246 (BOEHRINGER M.)

3/7


FIGURE 3



	Ra	Rb	Rc	Rd
1.	Cl	H	Cl	CH ₃
2.	H	H	OCF ₃	H
3.	H	H	CN	H
4.	H	H	SO ₂ CH ₃	H




	Ra	Rb	Rc	Rd
5.	H	H	CF ₃	H
6.	Cl	H	CF ₃	H
7.	H	H	COCH ₃	H
8.	H	Cl	Cl	H




	Ra	Rb	Rc	Rd
9.	Cl	H	Cl	H
10.	H	H	SCH ₃	H
11.	H	H	S(O)CH ₃	H

4/7


FIGURE 3 (continued)



	Ra	Rb	Rc	Rd
12.	Cl	H	Cl	H
13.	H	H	OCH ₃	H
14.	H	H	I	H
15.	H	H	C(O)CH ₃	H
16.	H	H	CF ₃	H




	Ra	Rb	Rc	Rd
17.	Cl	H	Cl	H
18.	H	H	CF ₃	H




	Ra	Rb	Rc	Rd
19.	Cl	H	Cl	H
20.	Cl	H	CF ₃	H
21.	H	H	I	H
22.	H	H	OCH ₃	H

5/7

FIGURE 3 (continued)




		Ra	Rb	Rc	Rd
23.	3,5-dichlorophenyl	Cl	H	Cl	H
24.	3,5-difluorophenyl	H	H	CF ₃	H
25.	3,5-dichlorophenyl	H	H	CF ₃	H
26.	3,5-difluorophenyl	Cl	H	Cl	H
27.	3,5-difluorophenyl	H	H	OCH ₃	H
28.	3,5-dimethoxyphenyl	Cl	H	Cl	H
29.	3,5-dimethoxyphenyl	Cl	H	CF ₃	H
30.	3,5-dimethoxyphenyl	H	H	OCH ₃	H




	Ra	Rb	Rc	Rd
31.	Cl	H	Cl	H
32.	Cl	H	I	H
33.	H	H	OCH ₃	H

6/7


FIGURE 3 (continued)



	Ra	Rb	Rc	Rd
34.	H	Cl	Cl	H
35.	Cl	H	Cl	H
36.	H	H	I	H




	Ra	Rb	Rc	Rd
37.	H	Cl	Cl	H
38.	Cl	H	Cl	H
39.	H	H	I	H
40.	Cl	H	Cl	Me
41.	Cl	H	CF ₃	H



	Ra	Rb	Rc	Rd	Re	Rf
42.	H	H	Cl	H	CH ₃	Cl
43.	H	H	Cl	H	CF ₃	Cl
44.	H	H	Cl	H	Br	Cl
45.	H	H	Cl	H	CF ₃	H

7/7

FIGURE 3 (continued)



	Ra	Rb	Rc	Rd
46.	H	Cl	Cl	H
47.	Cl	H	Cl	H
48.	H	H	I	H
49.	Cl	H	CF ₃	H

SF 1076409 v1